

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

BRAVECTO[®]

112.5 MG FLURALANER Chewable Tablets for Very Small Dogs
250 MG FLURALANER Chewable Tablets for Small Dogs
500 MG FLURALANER Chewable Tablets for Medium Dogs
1000 MG FLURALANER Chewable Tablets for Large Dogs
1400 MG FLURALANER Chewable Tablets for Very Large Dogs

ACTIVE CONSTITUENT: 136.4 g/kg Fluralaner

BRAVECTO Chewable Tablets provide

- Treatment and prevention of flea (*Ctenocephalides felis*) infestations for **3 months**
- Treatment and control of paralysis tick (*Ixodes holocyclus*) for **4 months**
- Treatment and control of brown dog tick (*Rhipicephalus sanguineus*) for **8 weeks**
- Rapid onset of action kills newly emerged adult fleas before they lay eggs
- Reduces incidence of flea allergy dermatitis (FAD) by controlling adult fleas
- Treatment and control of demodicosis (demodectic mange) caused by *Demodex* spp. mite infestations for up to **3 months**
- Treatment and control of sarcoptic mange (*Sarcoptes scabiei*) infestations for up to **1 month**
- Treatment and control of ear mite (*Otodectes cynotis*) infestations for up to **1 month**
- Highly palatable so easy to administer

BRAVECTO contains fluralaner, a member of the antiparasitic class of isoxazoline-substituted benzamide derivatives. Fluralaner acts antagonistically on arthropods ligand-gated chloride channels (GABA-receptor and glutamate-receptor). **BRAVECTO** is effective against fipronil-resistant strains of fleas.

BRAVECTO can be used in breeding, pregnant and lactating dogs.

ONSET OF ACTION AND EFFICACY

BRAVECTO provides effective control of pre-existing infestations of paralysis ticks (*Ixodes holocyclus*) within 24 hours.

179022 R4

BRAVECTO provides effective control of fleas on dogs within 8 hours of administration, and subsequently maintains persistent efficacy against flea re-infestations for **3 months** following administration. The flea life cycle can be broken by the rapid onset of action and the long-lasting efficacy against adult fleas on the dog. Newly emerged adult fleas on a dog are killed before viable eggs are produced.

BRAVECTO can be used as part of a treatment strategy for flea allergy dermatitis (FAD). *Ixodes holocyclus* ticks do not occur in Western Australia.

In tick season daily searching for and removal of any ticks found is recommended.

DIRECTIONS FOR USE

Precautions

BRAVECTO should not be used in puppies less than 8 weeks of age or weighing less than 2 kg.

Dosage and administration

BRAVECTO is available in five sizes. The following table indicates the size of chewable tablet to be used according to the bodyweight of the dog (equivalent to 25-56 mg fluralaner/kg bodyweight).

Bodyweight of dog (kg)	Chewable tablet size to be used
2 - 4.5	BRAVECTO 112.5 MG FLURALANER Chewable Tablets for Very Small Dogs
>4.5 - 10	BRAVECTO 250 MG FLURALANER Chewable Tablets for Small Dogs
>10 - 20	BRAVECTO 500 MG FLURALANER Chewable Tablets for Medium Dogs
>20 - 40	BRAVECTO 1000 MG FLURALANER Chewable Tablets for Large Dogs
>40 - 56	BRAVECTO 1400 MG FLURALANER Chewable Tablets for Very Large Dogs

Within each weight range, a whole tablet must be used. For dogs weighing more than 56 kg, use a combination of two tablets that most closely matches the bodyweight.

In dogs older than 8 weeks, treatment every 8 weeks has been shown to be well tolerated.

Puppies should be weighed regularly. Fast growing puppies that outgrow the initial weight band during the retreatment interval, may be retreated for fleas and paralysis ticks at 8 week intervals. Treatment may be appropriately adapted by the veterinarian to suit the individual weight changes.

Administer **BRAVECTO** at or around the time of feeding. **BRAVECTO** is readily consumed by most dogs when given by the owner.

BRAVECTO can also be given with food or administered in the same way as other tablets.

BRAVECTO chewable tablets can be used year round.

For further information contact MSD Animal Health Customer Service on 1800 033 461.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126.*

DISPOSAL

Dispose of empty foil and carton by wrapping in paper and putting in garbage.

STORAGE

Store below 30°C (room temperature).

APVMA:

BRAVECTO 112.5 MG FLURALANER Chewable Tablets for Very Small Dogs - 68867/107155

BRAVECTO 250 MG FLURALANER Chewable Tablets for Small Dogs - 68872/107154

BRAVECTO 500 MG FLURALANER Chewable Tablets for Medium Dogs - 68871/107153

BRAVECTO 1000 MG FLURALANER Chewable Tablets for Large Dogs - 68870/107152

BRAVECTO 1400 MG FLURALANER Chewable Tablets for Very Large Dogs - 68873/107150

Warranty

Intervet Australia Pty Limited (IAPL), trading as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.



Intervet Australia Pty Limited
(trading as MSD Animal Health)
91-105 Harpin Street,
Bendigo East VIC 3550
Phone: 1800 230 833
www.bravecto.com.au

® Registered Trademark